


Agarose



1% 1x TAE Universal Agarose gel showing separation of λ-DNA digested with EcoR I/Hind III [1] and a mixture from pBR328- DNA digested with Bgl I and Hinf I [2]. Data in kbp.

Features:

- High separation properties and sharp band patterns
- Easy solubility without foaming
- Excellent optical transparency

Applications:

- Separation of PCR products
 - DNA: approx. 0.05 kbp – 50 kbp
 - RNA: approx. 0.30 kb – 20 kb

Specifications:

- Gelling temperature: ≤ 37 °C
- Melting temperature: ≤ 90 °C
- Electroendosmosis: ≤ 0.140
- Gel strength (1.5 %): ≥ 2300 g / cm²
- Sulphate content: ≤ 0.10 %
- Water content: ≤ 10.0 %

Quality Assurance:

- 'Molecular Biology Grade'
- Certified free of DNases and RNases
- No DNA binding
- High lot-to-lot consistency

Usage:

Method 1: Microwave oven

1. Pour buffer (approx. 90 % of final volume) into an appropriate flask that can accommodate up to four times the final gel volume and add a magnetic stir bar.
2. Put the flask onto a magnetic stirrer and slowly add agarose powder while stirring constantly to avoid clotting.
3. Remove magnetic stir bar.
4. Add remaining buffer up to the desired final volume.
5. Weigh and record the weight of the flask prior to heating. Heat for 1 – 2 minutes in a microwave oven (600 Watt). Gently swirl the flask to mix the solution. Warning: Due to microwave heating, there may be a delay in the liquid boiling!
6. Using the microwave oven, heat in short bursts of 5 – 10 seconds or until the solution is boiling, with breaks of 10 – 15 seconds between heating phases to disperse bubbles by gently swirling the flask. Beware of hot glass ware and liquid. Continue until the agarose is completely dissolved.
7. Again, weigh the flask and top up lost volume with warm, deionized water. Gently swirl the flask to ensure complete mixing.
8. Let the solution cool down at room temperature for 15 – 20 minutes or until a gel temperature of 50 – 60 °C is reached.

Method 2: Simmering water bath

1. see method 1: steps 1 – 2 and 4
2. Weigh and record the weight of the flask prior to heating. Heat agarose suspension up in a simmering water bath with constant stirring.
3. Leave the flask in the water bath for further 15 – 20 minutes, or until the agarose is completely dissolved.
4. Switch off the magnetic stirrer and leave the flask in the bath for further 15 minutes.
5. Again, weigh the flask and top up lost volume with warm, deionized water. Gently swirl the flask to ensure complete mixing.
6. Let the solution cool down at room temperature for 15 – 20 minutes or until a gel temperature of 50 – 60 °C is reached.

Storage: room temperature for more than 4 years
Ordering information:

Cat.-no	Description	Amount
604-001	Agarose universal for gel-electrophoresis	100 g
604-005	Agarose universal for gel-electrophoresis	500 g

GeneON -Germany

Contact Phone +49-(0)-621- 5720 864 Fax: +49-(0)-621-5724 462
 E-Mail: <mailto:info@geneon.net> WEB: <http://www.GeneOn.net> Version: 10.2009 UK
 Unless specified otherwise, all products of GeneON are sold for research use only.

.. a good decision ..